

Lutein across the Lifespan: From Childhood Cognitive Performance to the Aging Eye and Brain

James M Stringham ¹, Elizabeth J Johnson ² and B Randy Hammond³

¹Visual performance Laboratory, Duke Eye Center, Durham, NC, USA; ²Friedman School of Nutrition, Tufts University, Boston, MA, USA; and ³Department of Psychology, University of Georgia-Athens, Athens, GA, USA

ABSTRACT

Lutein is a non-provitamin A dietary carotenoid found in dark green leafy vegetables, corn, eggs, and avocados. Among the carotenoids, lutein and its isomer, zeaxanthin, are the only 2 that cross the blood–retina barrier to form macular pigment in the retina. Lutein also preferentially accumulates in the human brain across multiple life stages. A variety of scientific evidence supports a role for lutein in visual as well as cognitive function across the lifespan. The purpose of this review is to summarize the latest science on lutein's role in the eye and the brain across different ages. *Curr Dev Nutr* 2019;3:nzz066.

Introduction

Lutein is a non-provitamin A dietary carotenoid found in dark green, leafy vegetables, corn, eggs, and avocados (1). Originally understood to preferentially accumulate in the macula of the eye, the majority of research characterized the effects of lutein and its isomer zeaxanthin on eye health. Lutein has been strongly implicated in visual function (2) as well as being protective against age-related eye diseases, particularly age-related macular degeneration (AMD) and cataracts (3, 4). More recently, insights into the bioavailability and bioaccumulation of lutein have led researchers to consider a role for lutein in brain development and cognitive performance. The purpose of this review is to summarize the latest science on lutein's role in the eye and the brain throughout the lifespan.

Effects of Lutein on Ocular Health and Visual Performance

The energy required for the countless physiological demands of the body is derived primarily from carbohydrates and lipids, composed of carbon and oxygen (4). In tissues with extremely high metabolism such as the retina (5) and brain (6), the instability of oxygen can cause devastating, often irreversible damage. Moreover, prolonged oxidative stress promotes proinflammatory responses (7), which can exacerbate damage and result in the creation of more reactive oxygen species (8). This positive feedback loop can result in the cumulative damage that is manifest in several age-related diseases, such as AMD (9, 10), atherosclerosis (11), and certain forms of cancer (12).

In the case of the retina, the metabolic rate is so exceedingly high that the body has prioritized a specific, nutrient-based strategy to help deal with the potential for oxidative stress. Namely, the carotenoids lutein and zeaxanthin accumulate in high densities in the region of the retina with the highest demand for oxygen, and therefore the highest risk of oxidative damage: the fovea (13). The fovea is extremely important to visual performance, because it mediates central vision and maintains the highest performance in terms of color discrimination, motion detection, contrast sensitivity, and acuity (14). Oxidative stress is not the only threat to the fovea, however—light



Keywords: lutein, zeaxanthin, carotenoids, nutrition, lifespan, eye, brain, vision, cognition

Copyright © American Society for Nutrition 2019. All rights reserved. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Manuscript received January 15, 2019. Initial review completed May 5, 2019. Revision accepted May 30, 2019. Published online June 4, 2019. This article was presented at the Nutrition 2018 meeting, Boston, MA, 9 June, 2018.

Supported by the Hass Avocado Board and the Egg Nutrition Center.

Author disclosures: EJJ was a scientific advisor for the Hass Avocado Board at the time of the writing of this manuscript. JMS, EJJ, and BRH were provided honorariums to present this topic at Nutrition 2018. There are no other conflicts of interest.

Publication costs for this supplement were defrayed in part by the payment of page charges by the Hass Avocado Board and the Egg Nutrition Center. The opinions expressed in this publication are those of the authors and are not attributable to the sponsors or the publisher, Editor, or Editorial Board of *Current Developments in Nutrition*.

Address correspondence to EJJ (e-mail: elizabeth.johnson@tufts.edu).

Abbreviations used: AMD, age-related macular degeneration; AREDS, Age-Related Eye Disease Study; CNS, central nervous system; IQ, intelligence quotient; MCI, mild cognitive impairment; MP, macular pigment; MPOD, macular pigment optical density; SES, social economic status.

energy incident on the retina can itself be damaging, whether via thermal- or photochemical-based lesions (15). This is especially true of short-wave visible light, which appears blue-violet in color. Owing to their yellow-orange coloration and location in the retina, lutein and zeaxanthin filter an appreciable amount of incident short-wave light before it reaches the photoreceptors of the sensory retina (16). Given their physical (light filtration) and biochemical (antioxidant) properties, lutein and zeaxanthin are ideally suited to protect the central retina and protect the lipid-dense retina from oxidation, and thereby facilitate good visual performance.

Lutein and retinal development

Despite the overwhelming focus on the association between relatively high levels of lutein in the diet/retina and a significant reduction in risk of developing AMD [e.g., (17, 18)], this relation is likely happenstance—adaptive pressures for accumulation of lutein in the retina would lead to survival advantages relatively early in life, and manifest before the reproductive cycle ends (not near the end of life). Indeed, there appears to be a significant role for lutein and zeaxanthin in the development of the retina. At the very earliest stages of retinal embryology (~14 weeks of gestation), lutein and zeaxanthin begin to accumulate in the large fluid reservoir that makes up most of the globular shape of the eye—the vitreous humor. This accumulation peaks at 20–22 weeks of gestation, at which point the lutein is diverted out and into the retinal space (19). Interestingly, this time period corresponds to the development of key layers of the retina such as Bruch's membrane, the plexiform layer, ganglion cell layer, nuclear layer, and photoreceptor layer (20). The timing suggests that lutein plays a role in the development of the retina. Additional evidence to support a role for lutein in retinal development comes from a study of premature infants (born <33 weeks of gestation) in which infant formula fortified with lutein, β -carotene, and lycopene was compared with a control formula with no added carotenoids (21). After ~4 mo of feeding with the respective formulas, electroretinography was performed to assess the status of retinal development. Overall, plasma lutein concentrations significantly correlated with the full-field electroretinogram response amplitude, and the supplemented group also showed greater rod photoreceptor sensitivity. In addition, the latency of light response was lower in the supplemented group, which indicates faster neural responses. These findings are suggestive of a significant and rapid contribution of lutein to retinal development. Lastly, it should be noted that lutein accumulates in the retina very early—perhaps in utero—at concentrations that are detectable with optical imaging techniques as early as 4 mo of age (22). The early and very specific accumulation in the retina further bolsters the argument for lutein as a key component of retinal development.

Lutein and visual performance

The presence of lutein and zeaxanthin in the central (macular) retina yields a yellow pigmentation, termed “macular pigment” (MP). The concentration of MP is typically expressed in terms of optical density (MPOD) and can range among individuals from 0 to values as high as 1.60 (23). This range represents a dramatic difference in both carotenoid concentration and light filtration: a value of 1.60 MPOD means that only ~2.5% of the short-wave visible light would penetrate through to the photoreceptors (compared with 100% for an individual with no MP). There have been several recent findings that indicate a significant

contribution by lutein and zeaxanthin to visual performance. Generally, visual performance has been shown to improve as a function of MPOD. This has been found to be true for a variety of performance parameters, including speed of visual processing, which is related linearly to MPOD (24, 25). This speed advantage also translates into accuracy, whereby positional judgments of quickly moving objects are more precise with higher MPOD (26). Another clear visual performance advantage of higher MPOD is visual performance in bright light conditions (i.e., glare). Visual discomfort has been found to be significantly inversely correlated with MPOD (27–29)—participants are able to tolerate more light before squinting or averting their gaze due to discomfort. In addition, glare disability (the phenomenon where glare produces “washed out” vision due to scattered light within the eye) is significantly reduced as a function of MPOD (27, 28, 30) and can also be significantly reduced within individuals who supplement with lutein and zeaxanthin (31–33). Similarly, photostress recovery (the time required to recover visibility after a relatively bright flash of light) is significantly shorter both with higher MPOD (30) and within individuals who augment their MPOD via dietary supplementation with lutein and zeaxanthin (33). Another aspect of visual performance that benefits from relatively high MPOD/augmentation of MPOD is contrast sensitivity. Contrast is the difference in luminance or color that makes objects distinguishable (34). The association between augmentation of MPOD and contrast sensitivity enhancement has been found in a number of recent studies in both normal individuals (35, 36) and an early-stage AMD clinical population (37). Improving visual performance in a progressive age-related disease such as AMD not only is clinically significant, but points to the plausibility of nutritional therapy to extend good visual function in diseased individuals, which could delay progression to the later, more debilitating stages. Lastly, visual function and performance in dim lighting conditions appear to benefit significantly from higher MPOD. This was first noted by Hammond et al. (38), in which dark-adapted visual sensitivity was significantly correlated with MPOD in older individuals (60–84 y). A similar finding in young, healthy participants was determined by Stringham et al. (39); they also found significantly enhanced speed of dark adaptation—the recovery time to see a very dim target after being in a bright environment was nearly 2 min less for individuals with high (compared with low) MPOD. This effect, however, may hold primarily for relatively young, healthy individuals. A recent report from Zarubina et al. (40) indicates that speed of dark adaptation is independent of MPOD in older individuals (>60 y).

Mechanisms of action

The aforementioned effects indicate the existence of several mechanisms of action for lutein and zeaxanthin in visual development and visual performance. The fact that these antioxidants confer protection to the retina—and thereby keep tissues healthy—accounts for at least some of these effects. In terms of visual performance in glare, short-wave light filtration would appear to be the most parsimonious explanation for performance enhancement. With regard to speed of visual processing and contrast sensitivity, light filtration cannot explain the significant enhancement seen in studies. Rather, these effects are most likely due to more efficient communication among visual neurons (41) and/or enhanced neurophysiological status in the retina and brain. Indeed, it was shown that MPOD is correlated significantly with a fundamental aspect of visual neurophysiology, called lateral inhibition (35). Lateral

inhibitory neural networks in the visual system ultimately lead to contrast sensitivity, and enhancement of contrast sensitivity by MPOD may be explained by the underlying optimization of neurophysiology, perhaps via promoting advantageous redox homeostasis (as suggested above).

It is important to note that the vast majority of Americans do not consume quantities of lutein and zeaxanthin sufficient to produce increases in MPOD (42). This is reflected in the relatively low averaged MPOD levels from 11 studies, conducted over a 15-y span, of ~ 0.35 (43). Given the significant improvements associated with MPOD increase (for both visual health and performance), the “room for improvement” is good news—dietary modification and/or supplementation with lutein and zeaxanthin can lead to relatively quick (<1 y), meaningful changes in visual health and performance.

Lutein and Cognition

Much of the early work on lutein and the central nervous system (CNS) focused on visual function/disease. This made sense because of the very high concentration of lutein and zeaxanthin in an area of the retina that is both critical to vision (cone-rich macula) and highly susceptible to disease (macular degeneration). It was not until relatively recently that some focus has turned to investigating the relation to other aspects of the CNS such as the brain. Recently, lutein has been observed to accumulate in the brain across multiple life stages ranging from infants to centenarians. Therefore, lutein may have an important role in the brain that is conserved across the lifespan. Lutein can be found in the circulation and throughout the human body where concentrations generally reflect intakes (44–48). Although lutein is not the major dietary carotenoid in the US diet for any age group (49), it is the carotenoid of highest concentration in human brain tissue (47, 50). This preferential accumulation in the brain is thought to occur owing to the presence of a specific lutein-binding protein present (51). In full-term human infants, lutein accounts for $\sim 60\%$ of total carotenoids in the brain despite survey data indicating lutein accounts for only $\sim 12\%$ of total carotenoids consumed in the diet in the first year of life (50). However, brain lutein content is significantly lower in preterm infants, who also have undetectable MPOD and very low concentrations of plasma carotenoids (52). Preferential accumulation of lutein in the brain is also observed at the opposite end of the lifespan, where lutein accounts for $\sim 35\%$ of total carotenoids in the centenarian brain despite making up only $\sim 20\%$ of the carotenoids in matched serum (47). Of interest is the suggestion that MPOD can serve as a biomarker of brain lutein concentrations. Certainly, this is plausible given that for lutein to be taken up in the retina it must cross the blood–brain barrier. Given this, MPOD measures might be a useful tool for cognitive studies testing the efficacy of lutein interventions. To test this, matched retina and brain sections from nonhuman primates (53) and humans (54) were analyzed for lutein. In nonhuman primates, lutein concentrations in the retina significantly correlated with lutein in the cerebellum, pons, and occipital cortex, and tended to correlate with concentrations in the frontal cortex. In humans, lutein in the retina was significantly related to lutein in the occipital cortex. These findings indicate that lutein accumulates up into the retina and multiple regions of the brain through similar mechanisms, thus supporting the use of MPOD as a

biomarker of lutein and its isomer, zeaxanthin, in human brain tissue. Of note, although lutein is the major dietary carotenoid in both retinal and brain tissues, concentrations were found to be considerably greater in the retina.

MP and cognitive function in the elderly

A number of recent studies have evaluated the relation between MPOD and cognitive function in adults. In the Health, Aging, and Body Composition Study, healthy older subjects ($n = 118$, 76–85 y) were assessed for MPOD and various measures of cognition. MPOD was significantly related to performance on a variety of indexes of processing speed, accuracy, and completion ability. These relations remained significant after adjusting for age, sex, and ethnicity. In the Irish Longitudinal Study on Aging ($n = 4453$, ≥ 50 y) (55), lower MPOD was significantly associated with poorer performance on the minimal state examination and on the Montreal cognitive assessment. Individuals with lower MPOD also had significantly poorer prospective memory, took significantly longer time to complete a trail-making task, and had significantly slower and more variable reaction times on a choice reaction time task. These associations were only slightly attenuated after adjustment for physical and mental health. Others investigating the relation between MPOD and cognitive function in healthy older adults (65–95 y, $n = 29$) and older adults with mild cognitive impairment (MCI) (65–95 y, $n = 24$) found that MPOD was associated with cognition in all subjects, but was more broadly correlated with cognitive test scores, including language, attention, and visual-spatial and constructional abilities, in MCI (56). More recently, Kelly et al. (57) conducted an investigation of the relation between MPOD and cognitive function in subjects free of retinal disease but low in MPOD (47 ± 12 y, $n = 105$) and subjects with AMD (65 ± 9 y, $n = 121$). Higher MPOD was significantly related to better performance in phonemic fluency, attention switching, visual and verbal memory, and learning even after adjusting for age, sex, diet, and education level. Collating across existing published studies, $>95\%$ of the subjects tested have been >50 y of age. This includes studies of centenarians (47), older subjects with mild cognitive decline (56, 58), and, most recently, patients with Alzheimer disease (23). Feeney et al. (59), for example, recently reported a positive relation between lutein status, global cognition, memory, and executive function in a sample of 4076 subjects over the age of 50 y. Collectively, these observations support the concept that MPOD reflects lutein content in brain tissue such that a relation between MPOD and cognitive function reflects lutein embedded in brain tissue.

Relation between brain concentrations of lutein and cognitive function

To further understand the relation between lutein status and cognitive function, the association between cognition and lutein concentrations in brain tissue of decedents >98 y at death and premortem measures of cognitive function were assessed. Subjects ($n = 29$) agreed to donate their brains upon death which were analyzed for carotenoids. Cognition measures included global cognition, primary degenerative dementia, delayed recall, delayed recognition, retention, intelligence quotient (IQ), and executive function (47). Among the carotenoids, brain lutein content was the most consistently related to cognitive test scores (47).

Lutein intervention and cognitive performance

Although the studies discussed to this point described compelling observational data on associations between lutein status and cognitive function in adults, the question remains, can lutein intake affect cognitive function? This question was examined in an intervention study involving a 4-mo, double-blind intervention trial of lutein and DHA (22:6n-3) supplementation (60). Forty-nine healthy women (60–80 y) were randomly assigned to receive lutein (12 mg/d) ($n = 11$), DHA (800 mg/d) ($n = 14$), a combination of lutein and DHA ($n = 14$), or placebo ($n = 10$). Cognitive tests including verbal fluency, memory, processing speed and accuracy, and self-reports of mood were performed at the study start and upon completion of the trial. After supplementation, verbal fluency scores improved significantly in the DHA-only, lutein-only, and combined treatment groups. Memory scores and rate of learning improved significantly in the lutein + DHA group, who also displayed a trend toward more efficient learning. This observation is consistent with findings from a study which reported a significant interaction between postmortem brain concentrations of lutein and DHA as a predictor of premortem cognitive test scores measuring verbal fluency, working memory, and dementia in centenarians (61). Combined lutein and DHA supplementation has also been shown to benefit younger adults (18–32 y old) (26). In this study, supplementation with lutein, zeaxanthin, and omega-3 fatty acids, including DHA, for 4 mo increased MP and improved neural processing speed. Collectively, these findings suggest that lutein may function in an additive/synergistic manner with DHA to influence cognitive function in older adults.

In a study involving older adults (mean age 63 y), a dietary intervention with avocados, a highly bioavailable food source of lutein, increased MPOD which was related to improved cognitive function (62). In other placebo-controlled intervention trials with older subjects (mean age 74 y), lutein/zeaxanthin supplements (12 mg/d for 1 y) resulted in cognitive improvements in older adults (mean age 74 y) (60, 63).

The influence of macular carotenoids on cognition in early life

Early work on lutein and the brain paralleled research on the visual system in that it tended to focus on older subjects and mechanisms that were mostly prophylactic. At first blush, it seems reasonable to infer that if lutein is influencing the cognitive function of subjects when they are older, it would also influence the cognition of subjects when they were younger. This inference is consistent with data showing that lutein/zeaxanthin supplementation improves the cognitive function of college students (64). Such generalizations, however, are not merited. Although much of the physical structure of the brain is complete by the age of 6 y (by some estimates, ~95%), modification of that structure continues until early adulthood. For instance, a second wave of synaptogenesis occurs near puberty (11–12 y) with significant pruning after that time (~1% of grey matter being lost every year between ~13–18 y) (65). The brain, like many biological systems, has a very close structure–function relation. Hence, significant maturational changes during adolescence mirror equally significant functional change. Because of such rapid change, the brain is particularly sensitive to exogenous input during adolescence. This has been shown very clearly by the extreme sensitivity of the adolescent brain to negative

factors such as stress (66). The reverse implication, however, is also true: the brain may have an equally exquisite sensitivity to positive factors such as a good diet (67).

Studies on the role of the macular carotenoids in the brain function of children are quite recent. The ability to investigate this relation has benefitted from the same advantage as the study of lutein in the brain of adults: namely, that retinal MP can be measured as a noninvasive biomarker for the amount of lutein in the brain (53, 54). The ability to measure a component of food, directly and nondestructively, in the very tissue that it could be affecting (the CNS) represents a powerful advantage. Concentrations of lutein when assessed in diet and serum are likely poor indicators of concentrations in the CNS. For example, Mulder et al. (48) measured lutein in the diet and serum of children in Canada and found that dietary intake (using the FFQ) only explained ~10% of the variation in serum lutein. This is similar to the type of correlations seen between serum lutein + zeaxanthin and MPOD measured in adults (correlations range around 0.30) (68). MPOD can, however, be reliably measured in preadolescent children (69) and, when used as a biomarker, provide more of a direct assessment of lutein's influence on CNS tissue.

Using this technology, 4 recent cross-sectional studies have assessed the relation between MPOD and cognitive function in preadolescent children. The first was by Hassevoort et al. (70). In that study, 40 children aged 7–10 y had their MPOD, adiposity, and aerobic fitness measured and compared with their performance on a relational memory task. This task is known to be mediated by the hippocampi which are also known to preferentially accumulate lutein in infancy (71). Hassevoort et al. found that MPOD was negatively related to relational memory errors. The correlations remained even after adjustments for IQ and other health factors such as $\dot{V}O_2$ max. The next study (72), using fifty-six 8- to 9-y-olds, studied academic achievement directly. Since passage of the No Child Left Behind Act in 2001, academic progress has increasingly focused on standardized testing of basic scholastic skills. The comprehensive version of the Kaufman Test of Academic and Educational Achievement was used to assess such basic skills as reading, spelling, and mathematics. MPOD related to achievement ($r = 0.40$, $P < 0.01$), reading ($r = 0.28$, $P < 0.05$), math ($r = 0.35$, $P < 0.01$), and written language ($r = 0.41$, $P < 0.01$). Saint et al. (73), testing fifty-one 7- to 13-y-olds, found similar-sized correlations using the Woodcock-Johnson III (composite standard scores were obtained for Brief Intellectual Ability, Verbal Ability, Cognitive Efficiency, Processing Speed, and Executive Processes): MPOD was significantly related to Executive Processes ($r = 0.29$, $P < 0.05$) and Brief Intellectual Ability ($r = 0.27$, $P < 0.05$). The studies on children appear to be reflecting those on adults; lutein measured directly tends to explain ~10% of the variance in a range of cognitive functions.

The results on children are unique in that they de-emphasize the prophylactic function of the pigments. In other words, it seems less likely that the brain function of the children is being improved by preventing degenerative change (through, for instance, an antioxidant or anti-inflammatory mechanism). The multiple-regression analyses by Hassevoort et al. (70) and Barnett et al. (72) further suggest that MPOD is not simply acting as a biomarker of improved health, IQ, or social economic status (SES). Walk et al. (74), using cognitive testing in conjunction with electroencephalography, and a matched case-control design (on variables like SES, age, and pubertal timing),

found that children with higher MPOD had lower brain activation when conducting the same cognitive tasks as children with lower MPOD. In other words, matched on a range of personal variables, children with lower MPOD had to use more brain to do the same task (decreased neural efficiency) and they made significantly more errors.

If these relations are not simply reflective of improved health, SES, and IQ, could MPOD be confounded with some other variables that relate to better cognition? Lieblein-Boff et al. (71) have shown that lutein relates to major amino acid neurotransmitters (like glutamate and γ -aminobutyric acid) that are well known to influence cognition. The fact that supplementing with lutein/zeaxanthin improves cognition in young adults, however, would suggest that, if this is the mechanism, then increasing lutein/zeaxanthin concentrations would also increase these neurotransmitters directly. This hypothesis can actually be tested because neurotransmitter levels in the brain can be measured noninvasively using available methods, e.g., Marsman et al. (75).

Mechanism of action

Despite consistent evidence of a beneficial relation of lutein with visual and cognitive function throughout the lifespan, the underlying mechanism of action remains unclear. It is speculated that some of the mechanisms by which lutein protects the retina from damage (antioxidant, anti-inflammatory) may also apply to the brain (76). The combination of high metabolic activity and rich PUFA content in the brain makes this organ vulnerable to oxidative stress (77). Furthermore, elevated DHA oxidation has been observed in patients with dementia and cognitive impairment (78). Unlike nonpolar carotenoids, such as β -carotene and lycopene, lutein has polar groups at each end of the molecule, which is believed to cause the molecule to span the membrane in a perpendicular or semiperpendicular orientation to the membrane surface (79, 80). This, in combination with evidence demonstrating that lutein localizes to membrane domains rich in PUFAs (including DHA) (81, 82), suggests that lutein is well positioned in membranes to block oxidation of these vulnerable brain lipids. Inhibition of DHA oxidation not only helps to maintain membrane structure and fluidity but also preserves DHA so it remains available for cleavage and conversion into anti-inflammatory molecules (83). Although this potential lipid-protective action by lutein may, in part, explain the relation between lutein and neural function, it is possible that lutein may function through other independent mechanisms, including modulation of membrane stability and function, as well as communications between neurons (79, 84).

Recommendations for Lutein Intake and Neural Health

Dietary sources

Rich food sources to consider are green leafy vegetables (spinach and kale), squash, broccoli, and corn (85, 86) (Table 1). Although eggs and avocados have relatively lower concentrations of lutein, both are considered highly bioavailable food sources of lutein, owing to their favorable fat profiles, and have been shown to increase MPOD and cognitive function in older adults (87–89).

TABLE 1 Lutein + zeaxanthin content in selected foods (85, 86)

Food	Serving	Lutein + zeaxanthin (mg)
Spinach, frozen, cooked	1 cup	29.8
Kale, frozen, cooked	1 cup	25.6
Summer squash, cooked	1 cup	4.0
Peas, frozen, cooked	1 cup	3.8
Pumpkin, cooked	1 cup	2.5
Brussel sprouts, frozen, cooked	1 cup	2.4
Broccoli, frozen, cooked	1 cup	2.0
Sweet yellow corn, boiled	1 cup	1.5
Avocado, raw	1 medium	0.4
Egg yolk, raw	1 large	0.2

Adults

It may be too early to make a recommendation for dietary intakes of lutein for promotion of cognitive health. However, intakes of ~ 6 mg/d have been related to a decreased risk of AMD (17), a disease which shares similar risk factors with age-related cognitive decline (90–92). Furthermore, the Age-Related Eye Disease Study 2 (AREDS2), a multicenter phase III randomized clinical trial, reported that in secondary analysis, lutein and zeaxanthin supplements (10 and 2 mg/d, respectively) on top of the AREDS supplement lowered the progression to advanced AMD in persons with low dietary lutein and zeaxanthin (93). And lastly, an intervention of 12 mg/d was found to improve cognition in older adults (60). Although these amounts (6–12 mg) can be achieved through proper dietary selections, the majority of the US adult population are not reaching these intakes (49, 94).

Infants and children

Similarly, it may be too soon to make recommendations for intakes of lutein in infants and children as it pertains to cognition. However, supplementation with lutein can increase lutein in breast milk and plasma of lactating mothers taking the supplement as well as corresponding plasma lutein concentrations in their infants (95). Furthermore, formulas supplemented with lutein enrich this bioactive in tissues, including retina and brain, compared with infants consuming lower amounts (96, 97). Therefore, higher intakes may be warranted to optimize brain development and cognition and recommendations for breastfeeding mothers may be necessary to ensure adequate lutein content in breast milk.

Dietary lutein appears to have an important role to play in the visual and cognitive development/optimization of children. If future intervention studies with children confirm these initial cross-sectional observations then they must also motivate the need to improve diet in early life. Lutein/zeaxanthin intake in American adults is low (~ 1 –2 mg/d) but it tends to be even lower in early adolescence (~ 300 –500 $\mu\text{g/d}$) (42). As with many aspects of diet, such deficits in childhood can start a cascade that simply amplifies with age. Interventions with children likely yield lifetime benefits.

Conclusions

To date, epidemiological evidence suggests that dietary lutein may be beneficial for visual and cognitive health across the lifespan. This

may be due to its role as an antioxidant and anti-inflammatory agent. Among the carotenoids, lutein preferentially accumulates into the infant and adult brain. A variety of evidence including observational and intervention studies suggests a relation between lutein status and visual and cognitive function in younger and older adults, as well as preadolescent children. Based on lutein intakes being related to a decreased risk of AMD, which has similar risk factors to age-related cognitive decline, average lutein intakes are low for adults. The elderly may be a particularly vulnerable group owing to poor nutrition for reasons including economics, medication use, and decreased sense of taste and smell. More recent observational evidence indicates that lutein is likely also important for brain development and cognition in early life. Breast milk is the optimal source of lutein for infants; however, lutein content in breast milk is dependent on maternal intake. Lutein-supplemented formulas may be used as an alternative source of lutein in infants.

Lutein is not an essential nutrient. However, evidence is accumulating to suggest that lutein is important to optimize ocular and cognitive health. Although the precise mechanisms by which lutein may be influencing neural health across the lifespan remain to be investigated, efforts may be warranted to establish recommended intakes for this dietary bioactive given the sum of the evidence to date and the low dietary intake of lutein across all age groups (49, 50).

Acknowledgments

All authors shared equal responsibility for the design and writing of this review. All authors have read and approved the final manuscript.

References

- Perry A, Rasmussen H, Johnson EJ. Xanthophyll (lutein, zeaxanthin) content in fruits, vegetables and corn and egg products. *J Food Compos Anal* 2009;22(1):9–15.
- Mares J. Lutein and zeaxanthin isomers in eye health and disease. *Annu Rev Nutr* 2016;36:571–602.
- Renzi LM, Johnson EJ. Lutein and age-related ocular disorders in the older adult: a review. *J Nutr Elder* 2007;26(3–4):139–57.
- Li W, Hu Z-F, Chen B, Ni G-X. Response of C2C12 myoblasts to hypoxia: the relative roles of glucose and oxygen in adaptive cellular metabolism. *BioMed Res Int* 2013;326346.
- Beatty S, Koh H, Phil M, Henson D, Boulton M. The role of oxidative stress in the pathogenesis of age-related macular degeneration. *Surv Ophthalmol* 2000;45(2):115–34.
- Floyd RA, Hensley K. Oxidative stress in brain aging. Implications for therapeutics of neurodegenerative diseases. *Neurobiol Aging* 2002;23(5):795–807.
- Nakanishi H, Wu Z. Microglia-aging: roles of microglial lysosome- and mitochondria-derived reactive oxygen species in brain aging. *Behav Brain Res* 2009;201(1):1–7.
- Afonso V, Santos G, Collin P, Khatib A-M, Mitrovic DR, Lomri N, Leitman DC, Lomri A. Tumor necrosis factor- α down-regulates human Cu/Zn superoxide dismutase 1 promoter via JNK/AP-1 signaling pathway. *Free Radic Biol Med* 2006;41(5):709–21.
- Liang F-Q, Godley BF. Oxidative stress-induced mitochondrial DNA damage in human retinal pigment epithelial cells: a possible mechanism for RPE aging and age-related macular degeneration. *Exp Eye Res* 2003;76(4):397–403.
- Lim LS, Mitchell P, Seddon JM, Holz FG, Wong TY. Age-related macular degeneration. *Lancet* 2012;379(9827):1728–38.
- Bonomini F, Tengattini S, Fabiano A, Bianchi R, Rezzani R. Atherosclerosis and oxidative stress. *Histol Histopathol* 2008;23(3):381–90.
- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 2006;160(1):1–40.
- Snodderly DM, Auran JD, Delori FC. The macular pigment. II. Spatial distribution in primate retinas. *Invest Ophthalmol Vis Sci* 1984;25(6):674–85.
- Schwiegerling J. Theoretical limits to visual performance. *Surv Ophthalmol* 2000;45(2):139–46.
- Ham WT, Mueller HA, Sliney DH. Retinal sensitivity to damage from short wavelength light. *Nature* 1976;260(5547):153–5.
- Snodderly DM, Brown PK, Delori FC, Auran JD. The macular pigment. I. Absorbance spectra, localization, and discrimination from other yellow pigments in primate retinas. *Invest Ophthalmol Vis Sci* 1984;25(6):660–73.
- Seddon JM, Ajani UA, Sperduto RD, Hiller R, Blair N, Burton TC, Farber MD, Gragoudas ES, Haller J, Miller DT, et al. Dietary carotenoids, vitamins A, C, and E, and advanced age-related macular degeneration. *JAMA* 1994;272(18):1413–20.
- Mares JA, Voland RP, Sondel SA, Millen AE, Larowe T, Moeller SM, Klein ML, Blodi BA, Chappell RJ, Tinker L, et al. Healthy lifestyles related to subsequent prevalence of age-related macular degeneration. *Arch Ophthalmol* 2011;129(4):470–80.
- Panova IG, Tatikolov AS, Sukhikh GT. Correlation between the content of albumin and carotenoids in human vitreous body during prenatal development. *Bull Exp Biol Med* 2007;144(5):681–3.
- Hendrickson A. Development of retinal layers in prenatal human retina. *Am J Ophthalmol* 2016;161:29–35. e1.
- Rubin LP, Chan GM, Barrett-Reis BM, Fulton AB, Hansen RM, Ashmeade TL, Oliver JS, Mackey AD, Dimmit RA, Hartmann EE, et al. Effect of carotenoid supplementation on plasma carotenoids, inflammation and visual development in preterm infants. *J Perinatol* 2012;32(6):418–24.
- Sharifzadeh M, Bernstein PS, Gellermann W. Reflection-based imaging of macular pigment distributions in infants and children. *J Biomed Opt* 2013;18(11):116001.
- Hammond BR Jr, Wooten BR, Snodderly DM. Individual variations in the spatial profile of human macular pigment. *J Opt Soc Am A Opt Image Sci Vis* 1997;14(6):1187–96.
- Hammond BR, Wooten BR. CFF thresholds: relation to macular pigment optical density. *Ophthalmic Physiol Opt* 2005;25(4):315–19.
- Stringham JM, Stringham NT, O'Brien KJ. Macular carotenoid supplementation improves visual performance, sleep quality, and adverse physical symptoms in those with high screen time exposure. *Foods* 2017;6(7):47.
- Bovier ER, Renzi LM, Hammond BR. A double-blind, placebo-controlled study on the effects of lutein and zeaxanthin on neural processing speed and efficiency. *PLoS One* 2014;9(9):e108178.
- Stringham JM, Fuld K, Wenzel AJ. Spatial properties of photophobia. *Invest Ophthalmol Vis Sci* 2004;45(10):3838–48.
- Stringham JM, Garcia PV, Smith PA, McLin LN, Foutch BK. Macular pigment and visual performance in glare: benefits for photostress recovery, disability glare, and visual discomfort. *Invest Ophthalmol Vis Sci* 2011;52(10):7406–15.
- Stringham JM, Snodderly DM. Enhancing performance while avoiding damage: a contribution of macular pigment. *Invest Ophthalmol Vis Sci* 2013;54(9):6298–306.
- Stringham JM, Hammond BR Jr. The glare hypothesis of macular pigment function. *Optom Vis Sci* 2007;84(9):859–64.
- Stringham JM, Hammond BR. Macular pigment and visual performance under glare conditions. *Optom Vis Sci* 2008;85(2):82–8.
- Hammond BR, Fletcher LM, Roos F, Wittwer J, Schalch W. A double-blind, placebo-controlled study on the effects of lutein and zeaxanthin on photostress recovery, glare disability, and chromatic contrast. *Invest Ophthalmol Vis Sci* 2014;55(12):8583–9.
- Stringham JM, O'Brien KJ, Stringham NT. Macular carotenoid supplementation improves disability glare performance and dynamics of photostress recovery. *Eye Vis* 2016;3:30.

34. Owsley C, Sloane ME. Contrast sensitivity, acuity, and the perception of “real-world” targets. *Br J Ophthalmol* 1987;71(10):791–6.
35. Stringham JM, O'Brien KJ, Stringham NT. Contrast sensitivity and lateral inhibition are enhanced with macular carotenoid supplementation. *Invest Ophthalmol Vis Sci* 2017;58(4):2291–5.
36. Nolan JM, Power R, Stringham J, Dennison J, Stack J, Kelly D, Moran R, Akuffo KO, Corcoran L, Beatty S. Enrichment of macular pigment enhances contrast sensitivity in subjects free of retinal disease: Central Retinal Enrichment Supplementation Trials – report 1. *Invest Ophthalmol Vis Sci* 2016;57(7):3429–39.
37. Akuffo KO, Beatty S, Peto T, Stack J, Stringham J, Kelly D, Leung I, Corcoran L, Nolan JM. The impact of supplemental antioxidants on visual function in nonadvanced age-related macular degeneration: a head-to-head randomized clinical trial. *Invest Ophthalmol Vis Sci* 2017;58(12):5347–60.
38. Hammond BR Jr, Wooten BR, Snodderly DM. Preservation of visual sensitivity of older subjects: association with macular pigment density. *Invest Ophthalmol Vis Sci* 1998;39(2):397–406.
39. Stringham JM, Garcia PV, Smith PA, Hiers PL, McLin LN, Kuyk TK, Foutch BK. Macular pigment and visual performance in low-light conditions. *Invest Ophthalmol Vis Sci* 2015;56(4):2459–68.
40. Zarubina AV, Huisingh CE, Clark ME, Sloan KR, McGwin G, Crosson JN, Curcio CA, Owsley C. Rod-mediated dark adaptation and macular pigment optical density in older adults with normal maculas. *Curr Eye Res* 2018;43(7):913–20.
41. Renzi LM, Hammond BR Jr. The relation between the macular carotenoids, lutein and zeaxanthin, and temporal vision. *Ophthalmic Physiol Opt* 2010;30(4):351–7.
42. Johnson EJ, Maras JE, Rasmussen HM, Tucker KL. Intake of lutein and zeaxanthin differ with age, sex, and ethnicity. *J Am Diet Assoc* 2010;110(9):1357–62.
43. Stringham NT, Stringham JM. Temporal visual mechanisms may mediate compensation for macular pigment. *Perception* 2015;44(12):1400–15.
44. Chung H-Y, Ferreira ALA, Epstein S, Paiva SAR, Castaneda-Sceppa C, Johnson EJ. Site-specific concentrations of carotenoids in adipose tissue: relations with dietary and serum carotenoid concentrations in healthy adults. *Am J Clin Nutr* 2009;90(3):533–9.
45. Hammond BR Jr, Johnson EJ, Russell RM, Krinsky NI, Yeum KJ, Edwards RB, Snodderly DM. Dietary modification of human macular pigment density. *Invest Ophthalmol Vis Sci* 1997;38(9):1795–801.
46. Mares JA, LaRowe TL, Snodderly DM, Moeller SM, Gruber MJ, Klein ML, Wooten BR, Johnson EJ, Chappell RJ; CAREDS Macular Pigment Study Group and Investigators. Predictors of optical density of lutein and zeaxanthin in retinas of older women in the Carotenoids in Age-Related Eye Disease Study, an ancillary study of the Women's Health Initiative. *Am J Clin Nutr* 2006;84(5):1107–22.
47. Johnson EJ, Vishwanathan R, Johnson MA, Hausman DB, Davey A, Scott TM, Green RC, Miller LS, Gearing M, Woodard J, et al. Relationship between serum and brain carotenoids, α -tocopherol, and retinol concentrations and cognitive performance in the oldest old from the Georgia Centenarian Study. *J Aging Res* 2013;95:1786.
48. Mulder KA, Innis SM, Rasmussen BF, Wu BT, Richardson KJ, Hasman D. Plasma lutein concentrations are related to dietary intake, but unrelated to dietary saturated fat or cognition in young children. *J Nutr Sci* 2014;3:e11.
49. Food and Nutrition Board, Institute of Medicine. Dietary Reference Intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. Washington (DC): National Academies Press; 2014.
50. Vishwanathan R, Kuchan MJ, Sen S, Johnson EJ. Lutein and preterm infants with decreased concentrations of brain carotenoids. *J Pediatr Gastroenterol Nutr* 2014;59(5):659–65.
51. Tanprasertsuk J, Li B, Bernstein PS, Vishwanathan R, Johnson MA, Poon L, Johnson EJ. Relationship between concentrations of lutein and StARD3 among pediatric and geriatric human brain tissue. *PLoS One* 2016;11(5):e0155488.
52. Bernstein PS, Sharifzadeh M, Liu A, Ermakov I, Nelson K, Sheng X, Panish C, Carlstrom B, Hoffman RO, Gellermann W. Blue-light reflectance imaging of macular pigment in infants and children. *Invest Ophthalmol Vis Sci* 2013;54(6):4034–40.
53. Vishwanathan R, Neuringer M, Snodderly DM, Schalch W, Johnson EJ. Macular lutein and zeaxanthin are related to brain lutein and zeaxanthin in primates. *Nutr Neurosci* 2013;16(1):21–9.
54. Vishwanathan R, Schalch W, Johnson EJ. Macular pigment carotenoids in the retina and occipital cortex are related in humans. *Nutr Neurosci* 2016;19(3):95–101.
55. Feeney J, Finucane C, Savva GM, Cronin H, Beatty S, Nolan JM, Kenny RA. Low macular pigment optical density is associated with lower cognitive performance in a large, population-based sample of older adults. *Neurobiol Aging* 2013;34(11):2449–56.
56. Renzi LM, Dengler MJ, Puente A, Miller LS, Hammond BR. Relationships between macular pigment optical density and cognitive function in unimpaired and mildly cognitively impaired older adults. *Neurobiol Aging* 2014;35(7):1695–9.
57. Kelly D, Coen RF, Akuffo KO, Beatty S, Dennison J, Moran R, Stack J, Howard AN, Mulcahy R, Nolan JM. Cognitive function and its relationship with macular pigment optical density and serum concentrations of its constituent carotenoids. *J Alzheimers Dis* 2015;48(1):261–77.
58. Nolan JM, Mulcahy R, Power R, Moran R, Howard AN. Nutritional intervention to prevent Alzheimer's disease: potential benefits of xanthophyll carotenoids and omega-3 fatty acids combined. *J Alzheimers Dis* 2018;64(2):367–78.
59. Feeney J, O'Sullivan M, Kenny RA, Robertson IH. Change in perceived stress and 2-year change in cognitive function among older adults: the Irish Longitudinal Study on Ageing. *Stress Health* 2018;34(3):403–10.
60. Johnson EJ, McDonald K, Caldarella SM, Chung H-Y, Troen AM, Snodderly DM. Cognitive findings of an exploratory trial of docosahexaenoic acid and lutein supplementation in older women. *Nutr Neurosci* 2008;11(2):75–83.
61. Mohn E, Vishwanathan R, Schalch W, Lichtenstein AH, Matthan NR, Poon LW, Green RC, Gearing M, Johnson EJ. The relationship of lutein and DHA in age-related cognitive function. Poster Presentation at Experimental Biology Conference; 20–24 April, 2013; Boston, MA.
62. Scott TM, Rasmussen HM, Chen O, Johnson EJ. Avocado consumption increases macular pigment density in older adults: a randomized, controlled trial. *Nutrients* 2017;9(9):E919.
63. Hammond BR, Miller LS, Bello MO, Lindbergh CA, Mewborn C, Renzi-Hammond LM. Effects of lutein/zeaxanthin supplementation on the cognitive function of community dwelling older adults: a randomized, double-masked, placebo-controlled trial. *Front Aging Neurosci* 2017;9:254.
64. Renzi-Hammond LM, Bovier ER, Fletcher LM, Miller LS, Mewborn CM, Lindbergh CA, Baxter JH, Hammond BR. Effects of a lutein and zeaxanthin intervention on cognitive function: a randomized, double-masked, placebo-controlled trial of younger healthy adults. *Nutrients* 2017;9(11):E1246.
65. Giedd JN, Raznahan A, Alexander-Bloch A, Schmitt E, Gogtay N, Rapoport JL. Child psychiatry branch of the National Institute of Mental Health longitudinal structural magnetic resonance imaging study of human brain development. *Neuropsychopharmacology* 2015;40(1):43–9.
66. Romeo RD. The metamorphosis of adolescent hormonal stress reactivity: a focus on animal models. *Front Neuroendocrinol* 2018;49:43–51.
67. Deoni SCL. Neuroimaging of the developing brain and impact of nutrition. *Nestle Nutr Inst Workshop Ser* 2018;89:155–74.
68. Ciulla TA, Curran-Celantano J, Cooper BR, Hammond BR Jr, Danis RP, Pratt LM, Riccardi KA, Filloon TG. Macular pigment optical density in a midwestern sample. *Ophthalmology* 2001;108(4):730–7.
69. McCorkle SM, Raine LB, Hammond BR Jr, Renzi-Hammond L, Hillman CH, Khan NA. Reliability of heterochromatic flicker photometry in measuring macular pigment optical density among preadolescent children. *Foods* 2015;4(4):594–604.
70. Hassevoort KM, Khazoum SE, Walker JA, Barnett SM, Raine LB, Hammond BR, Renzi-Hammond LM, Kramer AF, Khan NA, Hillman CH, et al. Macular carotenoids, aerobic fitness, and central adiposity are associated differentially with hippocampal-dependent relational memory in preadolescent children. *J Pediatr* 2017;183:108–14. e1.

71. Lieblein-Boff JC, Johnson EJ, Kennedy AD, Lai C-S, Kuchan MJ. Exploratory metabolomic analyses reveal compounds correlated with lutein concentration in frontal cortex, hippocampus, and occipital cortex of human infant brain. *PLoS One* 2015;10(8):e0136904.
72. Barnett SM, Khan NA, Walk AM, Raine LB, Moulton C, Cohen NJ, Kramer AF, Hammond BR Jr, Renzi-Hammond L, Hillman CH. Macular pigment optical density is positively associated with academic performance among preadolescent children. *Nutr Neurosci* 2018;21(9):632–40.
73. Saint SE, Renzi-Hammond LM, Khan NA, Hillman CH, Frick JE, Hammond BR. The macular carotenoids are associated with cognitive function in preadolescent children. *Nutrients* 2018;10(2):E193.
74. Walk AM, Khan NA, Barnett SM, Raine LB, Kramer AF, Cohen NJ, Moulton CJ, Renzi-Hammond LM, Hammond BR, Hillman CH. From neuro-pigments to neural efficiency: the relationship between retinal carotenoids and behavioral and neuroelectric indices of cognitive control in childhood. *Int J Psychophysiol* 2017;118:1–8.
75. Marsman A, Egerton A, Broberg B, Poh H. MR spectroscopy in neuropsychiatry. Lausanne, Switzerland: Frontiers Media SA; 2018.
76. Erdman JW Jr, Smith JW, Kuchan MJ, Mohn ES, Johnson EJ, Rubakhin SS, Wang L, Sweedler JV, Neuringer M. Lutein and brain function. *Foods* 2015;4(4):547–64.
77. Friedman J. Why is the nervous system vulnerable to oxidative stress? In: Gadoth N Göbel HH, editors. *Oxidative stress and free radical damage in neurology* [Internet]. New York: Humana Press; 2011 [cited 4 May, 2017]. p. 19–27. *Oxidative Stress in Applied Basic Research and Clinical Practice*. Available from: http://link.springer.com/chapter/10.1007/978-1-60327-514-9_2.
78. Reich EE, Markesbery WR, Roberts LJ, Swift LL, Morrow JD, Montine TJ. Brain regional quantification of F-ring and D-/E-ring isoprostanes and neuroprostanes in Alzheimer's disease. *Am J Pathol* 2001;158(1):293–7.
79. Krinsky NI, Mayne ST, Sies H. *Carotenoids in health and disease*. Boca Raton, FL: CRC Press; 2004.
80. Gruszecki W, Sujak A, Strzalka K, Radunz A, Schmid G. Organisation of xanthophyll-lipid membranes studied by means of specific pigment antisera, spectrophotometry and monomolecular layer technique lutein versus zeaxanthin. *Z Naturforschung C J Biosci* 1999;54(7–8):517–25.
81. Rapp LM, Maple SS, Choi JH. Lutein and zeaxanthin concentrations in rod outer segment membranes from perifoveal and peripheral human retina. *Invest Ophthalmol Vis Sci* 2000;41(5):1200–9.
82. Wisniewska A, Subczynski WK. Accumulation of macular xanthophylls in unsaturated membrane domains. *Free Radic Biol Med* 2006;40(10):1820–6.
83. Farooqui AA, Horrocks LA, Farooqui T. Modulation of inflammation in brain: a matter of fat. *J Neurochem* 2007;101(3):577–99.
84. Stahl W, Sies H. Effects of carotenoids and retinoids on gap junctional communication. *BioFactors* 2001;15(2–4):95–8.
85. US Department of Agriculture, Agricultural Research Service. USDA National Nutrient Database for Standard Reference, release 28 [Internet]. 2015. Available from: <http://ars.usda.gov/Services/docs.htm?docid=8964>.
86. Unlu NZ, Bohn T, Clinton SK, Schwartz SJ. Carotenoid absorption from salad and salsa by humans is enhanced by the addition of avocado or avocado oil. *J Nutr* 2005;135(3):431–6.
87. Chung H-Y, Rasmussen HM, Johnson EJ. Lutein bioavailability is higher from lutein-enriched eggs than from supplements and spinach in men. *J Nutr* 2004;134(8):1887–93.
88. Johnson E, Vishwanathan R, Mohn E, Haddock J, Rasmussen H, Scott T. Avocado consumption increases neural lutein and improves cognitive function. *FASEB J* 2015;29(1 Supplement):32.8.
89. Wenzel AJ, Gerweck C, Barbato D, Nicolosi RJ, Handelsman GJ, Curran-Celentano J. A 12-wk egg intervention increases serum zeaxanthin and macular pigment optical density in women. *J Nutr* 2006;136(10):2568–73.
90. Woo SJ, Park KH, Ahn J, Choe JY, Jeong H, Han JW, Kim TH, Kim KW. Cognitive impairment in age-related macular degeneration and geographic atrophy. *Ophthalmology* 2012;119(10):2094–101.
91. Wong TY, Klein R, Nieto FJ, Moraes SAD, Mosley TH, Couper DJ, Klein BE, Boland LL, Hubbard LD, Sharrett AR. Is early age-related maculopathy related to cognitive function? The Atherosclerosis Risk in Communities Study. *Am J Ophthalmol* 2002;134(6):828–35.
92. Pham TQ, Kifley A, Mitchell P, Wang JJ. Relation of age-related macular degeneration and cognitive impairment in an older population. *Gerontology* 2006;52(6):353–8.
93. Age-Related Eye Disease Study 2 Research Group. Lutein + zeaxanthin and omega-3 fatty acids for age-related macular degeneration: the Age-Related Eye Disease Study 2 (AREDS2) randomized clinical trial. *JAMA* 2013;309(19):2005–15.
94. Rasmussen HM, Johnson EJ. Nutrients for the aging eye. *Clin Interv Aging* 2013;8:741–8.
95. Sherry CL, Oliver JS, Renzi LM, Marriage BJ. Lutein supplementation increases breast milk and plasma lutein concentrations in lactating women and infant plasma concentrations but does not affect other carotenoids. *J Nutr* 2014;144(8):1256–63.
96. Jeon S, Neuringer M, Johnson EE, Kuchan MJ, Pereira SL, Johnson EJ, Erdman JW. Effect of carotenoid supplemented formula on carotenoid bioaccumulation in tissues of infant rhesus macaques: a pilot study focused on lutein. *Nutrients* 2017;9(1):E51.
97. Bettler J, Zimmer JP, Neuringer M, DeRusso PA. Serum lutein concentrations in healthy term infants fed human milk or infant formula with lutein. *Eur J Nutr* 2010;49(1):45–51.